



Final Baseline Human Health and Ecological Risk Assessment

Lower Fox River and Green Bay, Wisconsin Remedial Investigation and Feasibility Study

Prepared for:

Wisconsin Dept. of Natural Resources



Prepared by: The RETEC Group, Inc.

December 2002

Final Baseline Human Health and Ecological Risk Assessment

Lower Fox River and Green Bay, Wisconsin Remedial Investigation and Feasibility Study

Prepared by:

**The RETEC Group, Inc.
1011 S.W. Klickitat Way, Suite #207
Seattle, Washington 98134**

and

**3040 William Pitt Way
Pittsburgh, Pennsylvania 15238**

RETEC Project No.: WISCN-14414-461

Prepared for:

**Wisconsin Department of Natural Resources
101 S. Webster Street
Madison, Wisconsin 55703**

Prepared by:

**Timothy A. Thompson, Senior Environmental Scientist
David Morgan, Senior Technical Consultant
Linda Mortensen, Environmental Scientist
Annette Pearson, Environmental Scientist
Damon Morris, Environmental Scientist
Chris Traynor, Project Scientist**

December 2002

EXECUTIVE SUMMARY

A *Baseline Human Health and Ecological Risk Assessment for the Lower Fox River and Green Bay* (BLRA) has been prepared as a companion document to the Remedial Investigation (RI) and Feasibility Study (FS). This section summarizes the baseline risks to human health for the Lower Fox River and Green Bay, and the calculation of sediment quality thresholds (SQTs) that support the selection of a remedy which eliminates, reduces, and/or controls risks identified in the human health and ecological assessments. The SQTs themselves are not cleanup criteria, but are a good approximation of protective sediment values and can be considered to be “working values” from which to select a remedial action level.

This RI/FS report is consistent with the findings of the National Academy of Science’s National Research Council Report entitled, *A Risk Management Strategy for PCB Contaminated Sediments* (NRC, 2001).

The overall goals of the BLRA for the Lower Fox River and Green Bay were to:

- Examine how the contaminants of potential concern (COPCs) carried forward from the Screening Level Risk Assessment (SLRA) (RETEC, 1998b) move from the sediment and water into human and ecological receptors within the Lower Fox River and Green Bay.
- Quantify the current (or baseline) human health and ecological risk associated with the COPCs.

- Distinguish those COPCs which pose the greatest potential for risk to human health and the environment and should be carried forward as contaminants of concern (COCs) in the FS.
- Determine which exposure pathways lead to the greatest risks.
- Support the selection of a remedy which eliminates, reduces, and/or controls identified risks by calculating sediment quality thresholds (SQTs).




Figure 1 Risk Assessment Study Areas

Site Description

Between 1954 and 1971, paper mills in the Lower Fox River valley manufactured and recycled carbonless copy paper that contained PCBs, resulting in the release of an estimated 313,600 kg (691,370 pounds) of PCBs in the river. It is estimated that 70 percent of the total PCB mass in the river has been transported into Green Bay. Sediment from the Lower Fox River is primarily deposited on the southeastern edge of the bay.

The Fox River valley and Green Bay area is diverse in terms of land use, population density, and habitat types. Overall, the shoreline is much more developed and populated along the Lower Fox River as compared to Green Bay. Both the human health and ecological risk assessments focused on aquatic-dependent receptors and Green Bay has historically supported strong commercial and sport fishing.

For both the human health and ecological assessments, risk was characterized for the four reaches of the Lower Fox River: Little Lake Butte des Morts, Appleton to Little Rapids, Little Rapids to De Pere, and De Pere to Green Bay (Green Bay Zone 1); as well as the zones of the bay: Zone 2, Zone 3A, Zone 3B, and Zone 4 (Figure 1). Therefore, risks between each of these reaches and zones could be compared.

Data Evaluated

The COPCs carried forward from the SLRA included polychlorinated biphenyls (PCBs) (total and selected congeners), dioxins and furan congeners, dichlorodiphenyltrichloroethane (DDT) and its metabolites (4,4'-dichlorodiphenyltrichloroethane [DDE] and 4,4'-dichlorodiphenyldichloroethane [DDD]), dieldrin, and three metals (arsenic, lead, and mercury). In the SLRA, hazard quotients (HQs) calculated for PCBs were at least an order of magnitude greater than the HQs for any of the other COPCs. HQs are the ratios of measured COPC concentrations in media (water, sediment, tissue) as compared to safe COPC concentrations in these media.

All available electronic data collected from Lake Winnebago to northern Green Bay were compiled into a single database—the Fox River Database (FRDB). This database

contains 474,218 records of sediment, water, and tissue data from the early 1970s through the late 1990s. For the assessment of baseline risk in the Lower Fox River and Green Bay, a subset of the data contained in the FRDB was evaluated. Data were included based on the specific receptors selected, the time during which the data were collected, and the COPCs of interest.

A time trend analysis of fish tissue data indicates that while PCB concentrations in fish tissue initially significantly decreased, since the mid 1980s changes in these concentrations have either slowed, remained constant, or have resulted in increased tissue concentrations. For this reason, only fish tissue concentrations from 1989 and after were considered for the ecological risk evaluation and the focused human health risk evaluation.

Similarly, for risk evaluation purposes, the concentration of total PCBs in the top 10 cm (4 inches) of sediment was interpolated, because this is the depth of sediment that is of primary biological activity. The degree of biological activity influences the potential for bioaccumulative compounds to be taken up in the food chain. PCB concentrations in sediment were interpolated both horizontally and vertically, but for comparative risk purposes non-interpolated sediment PCB concentrations were also evaluated for risk.

General Conclusions

General conclusions of both the human health and ecological assessments were that:

- Fish consumption is the exposure pathway that represents the greatest level of risk for receptors (other than direct risk to benthic invertebrates).

- The primary COC is PCBs, and other COCs carried forward for remedial evaluation and long-term monitoring are mercury and DDE.
- In general, areas evaluated with the greatest risk are Green Bay zones 1 and 2.

Human Health Risk Assessment

For the human health risk assessment, two evaluations were performed, a baseline risk assessment and a focused risk assessment, which are described shortly. For the baseline risk assessment, all data for a specific medium for each COC were used to evaluate exposures and risks. For the focused risk assessment, which examined only exposure to PCBs in fish, only fish tissue data from 1989 and after were used.

Receptors evaluated in the human health risk assessment were:

- Recreational anglers,
- High-intake fish consumers,
- Hunters,
- Drinking water users,
- Local residents,
- Recreational water users (swimmers and waders), and
- Marine construction workers.

The principle findings of the human health risk assessment are:

- Consumption of fish from the Lower Fox River and Green Bay presented the

highest cancer risks and noncancer hazard indices for the pathways evaluated which also included those associated with consumption of waterfowl, drinking water, breathing air near the river or bay, swimming, and construction in the river or bay.

- PCBs contribute more than 70 percent of the cancer risks found from the consumption of fish and waterfowl.
- Using fish data since 1989, lifetime cancer risks as great as one in 1,000 were found for recreational anglers and high-intake fish consumers exposed to PCBs. High-intake fish consumers are individuals in the recreational angler population who may eat significantly more fish than recreational anglers. Groups within the high-intake fish consumer category that were explicitly evaluated in this risk assessment were low-income minority anglers, Hmong/Laotian anglers, and Native American anglers.
- While high-intake fish consumers are individuals who may eat significantly more fish than typical recreational anglers, there were not large differences in risks between recreational anglers and high-intake fish consumers for the high fish consumption or reasonable maximum exposure scenarios.
- Cancer risks from fish consumption are 1,000 times greater than the one-in-a-million cancer risk, which is the point at which risk management decisions may be made under Superfund. The cancer risks are 100 times greater than the one-in-a-hundred-thousand lifetime cancer risk

used by Wisconsin for evaluating hazardous waste sites.

- Noncancer hazard indices from fish consumption were as much as 50 times greater than levels considered acceptable for exposures ranging from 7 years to a lifetime. The noncancer health effects

density. The hazard indices were approximately 2.4 times those found for adults or as much as 125 times greater than acceptable levels.

- Populations potentially exposed to PCBs via fish consumption are large. There are 136,000 fishing licenses issued to

Table 1 Summary of Human Health Risks

Location	Recreational Angler		High Intake Fish Consumer		Hunter		Drinking Water User		Local Resident		Swimmer		Wader		Marine Construction Worker	
	Cancer Risk	Noncancer Hazard Index	Cancer Risk	Noncancer Hazard Index	Cancer Risk	Noncancer Hazard Index	Cancer Risk	Noncancer Hazard Index	Cancer Risk	Noncancer Hazard Index	Cancer Risk	Noncancer Hazard Index	Cancer Risk	Noncancer Hazard Index	Cancer Risk	Noncancer Hazard Index
Little Lake Butte des Morts	$>10^{-4}$	>25	$>10^{-4}$	>35	10^{-6} – 10^{-4}	~ 1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$\sim 10^{-6}$	<1
Appleton to Little Rapids	$>10^{-4}$	>20	$>10^{-4}$	>30	10^{-6} – 10^{-4}	~ 1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1
Little Rapids to De Pere	$>10^{-4}$	>15	$>10^{-4}$	>20	10^{-6} – 10^{-4}	~ 1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1
De Pere to Green Bay	$>10^{-4}$	>35	$>10^{-4}$	>50	10^{-6} – 10^{-4}	~ 1	10^{-6} – 10^{-4}	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1
Zone 3A	$>10^{-4}$	>25	$>10^{-4}$	>50	10^{-6} – 10^{-4}	~ 1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1	$<10^{-6}$	<1
Zone 3B	$>10^{-4}$	>25	$>10^{-4}$	>35												
Zone 4	$>10^{-4}$	>25	$>10^{-4}$	>35												

Notes:

Risks and hazard indices are based on reasonable maximum exposures.

Interpretation of cancer risks:

$>10^{-4}$ indicates significant risk

10^{-6} – 10^{-4} indicates possibly significant risks

$<10^{-6}$ indicates risks are negligible

Interpretation of hazard indices:

>1 indicates significant noncancer health effects are possible

<1 indicates noncancer health effects are unlikely to be significant

For recreational anglers, high intake fish consumers and hunters, elevated risks and hazard indices are due primarily to PCBs.

For drinking water users in De Pere to Green Bay reach, arsenic is responsible for calculated cancer risk, but arsenic exposure point concentration was based on one detected value in four samples and reporting limits were high, so actual arsenic concentration

associated with exposure to PCBs include developmental effects (e.g., neurological impairment in infants and children due to maternal exposure), reproductive effects (e.g., conceptive failure), and immune system suppression (e.g., increased incidence of infectious disease in infants).

- Noncancer hazard indices were also calculated for young children eating fish for the Little Lake Butte des Morts and De Pere to Green Bay reaches, the two reaches with the greatest population

individuals living in counties adjacent to the Lower Fox River and Green Bay. About 10 percent of this angler population, or about 14,000 persons, would be considered high-intake anglers. These populations are potentially exposed to PCBs at levels associated with adverse health consequences.

- Cancer risks and noncancer hazard indices are more than 20 times greater than those from the consumption of fish from Lake Winnebago, which does not

have a known source of PCBs and serves as a background location.

- There were not large differences in risks between the Lower Fox River and Green Bay, or among the reaches within the Lower Fox River, or among the zones within Green Bay.
- While evidence exists for slow declines of PCBs in fish, such declines were not consistent among species or locations, and projections of future declines cannot be made with sufficient certainty for use in risk assessment. In addition, in some cases, PCBs were found to be increasing.


Other findings of the human health risk assessment are:

- Cancer risks to hunters consuming waterfowl approach a risk of one in 10,000. Noncancer hazard indices were 3.8 times acceptable levels.
- Cancer risks to local residents exposed to chemicals only through inhalation of air, swimmers, and waders were less than one in a million.
- Cancer risks to drinking water users were less than one in a million in all reaches of the Lower Fox River and all of Green Bay with one exception. The cancer risk in the De Pere to Green Bay Reach was 3.8×10^{-5} due to exposure to arsenic. The arsenic and the exposure to arsenic were based on the detection of this chemical in one of four surface water samples. It is quite likely that this one detected value is anomalous and that the actual risk of exposure to arsenic is much lower. In addition, this reach of the

Lower Fox River is not used as a source of drinking water.

- Marine construction workers had cancer risks slightly greater than one in a million. Noncancer hazard indices for drinking water users, local residents, swimmers, waders, and marine construction workers did not exceed acceptable levels.

These results are summarized in Table 1. Figure 2 presents the risks and Figure 3 presents the hazard indices for recreational anglers and high-intake fish consumers due to ingestion of PCBs in fish.



Ecological Risk Assessment

Types of receptors evaluated for ecological risk included:


- **Aquatic Invertebrates:** Insects and other invertebrates that live in the water and are important prey items for fish and other insects.
- **Benthic Invertebrates:** Insects and other invertebrates that live in or on the sediment that are important in recycling

nutrients and are a principal part of fish diets.

- **Benthic Fish:** Fish, such as carp and catfish, that live on and forage in the sediments and are in turn eaten by other fish, birds, mammals, and people.
- **Pelagic Fish:** Fish, such as walleye and yellow perch, that live in the water column, and eat other fish or insects that live in the water or on the sediments. These fish may be in turn eaten by other fish, birds, mammals, and people.
- **Insectivorous Birds:** Birds, such as swallows, that eat insects that hatch from the sediments.
- **Piscivorous Birds:** Birds, such as cormorants or terns, that principally eat fish from the Lower Fox River or Green Bay.
- **Carnivorous Birds:** Birds, such as eagles, that eat a variety of prey, including fish or small mammals.
- **Piscivorous Mammals:** Mammals, such as mink, that eat fish as an important part of their diet.

Risk was characterized for assessment endpoints based on the calculation of HQs. In the FRDB, data were generally lacking for piscivorous and carnivorous birds, and no data were available for piscivorous mammals, therefore, ecological modeling was used to estimate COPC exposure to these receptors. HQs that are greater than 1.0 imply that risk may be present. Where available, both the No Observed Adverse Effect Concentration (NOAEC) and Lowest Observed Adverse Effect Concentration (LOAEC) HQs were

calculated. Effects evaluated were reproductive dysfunction, death at birth, or deformities in the surviving offspring. When NOAEC HQs exceeded 1.0, but LOAEC HQs were less than 1.0, then it was concluded that there was potential risk. When both the NOAEC and LOAEC HQs exceed 1.0, it was assumed that risk is present.



In addition to the HQ, the assessment provides an evaluation of the uncertainties associated with the risk characterization, and evaluates the estimated risk relative to the habitat, field studies, and population data for the receptors species. Together with the HQs, the components of the evaluation provide resource managers with the information necessary to make risk decisions within the context of the Feasibility Study.

The principle findings of the ecological risk assessment are:

- Total PCBs cause, or potentially cause risk to all identified receptors. The exception is insectivorous birds where the weight of evidence suggests that these receptors are not at risk from PCB

concentrations. Not all receptors at risk or potentially at risk from PCBs are at risk in all river reaches or bay zones.

- Mercury poses a risk in all river reaches and zones, but not to all receptors. Mercury was not identified as a risk for benthic fish, insectivorous birds, or piscivorous mammals.
- DDT or its metabolites poses a risk to benthic invertebrates (Little Lake Butte des Morts Reach, Little Rapids to De Pere Reach, and Green Bay Zone 1), benthic fish (Green Bay zones 1 and 2), pelagic fish (Green Bay zones 1, 2, 3B, and 4), insectivorous birds (Green Bay Zone 2), piscivorous birds (Green Bay zones 1, 2,

- Other COPCs identified as causing or potentially causing risk are arsenic (Zone 1 and Zone 3B benthic invertebrates only) lead (benthic invertebrates only in all areas except Green Bay Zone 2, Zone 3A, and Zone 4), 2,3,7,8-TCDD (benthic invertebrates only in Little Lake Butte des Morts Reach and Little Rapids to De Pere Reach), and dieldrin (piscivorous birds in zones 1, 2, and 3B, carnivorous birds in Green Bay Zone 3A, and piscivorous mammals in Green Bay zones 3A and 3B).

Table 2 summarizes ecological risks based on hazard quotients and other lines of evidence. Figures 4 (total PCBs), 5 (mercury), and 6

Table 2 Ecological Risk Summary Table

Location	Water Column Invertebrates	Benthic Invertebrates	Benthic Fish	Pelagic Fish	Insectivorous Bird	Piscivorous Bird	Carnivorous Bird	Piscivorous Mammal
LLBdM	● mercury ○ PCBs	● lead; mercury; 2,3,7,8-TCDD; PCBs; DDD; DDT	○ PCBs	○ PCBs	○ PCBs	○ mercury; PCBs	○ PCBs	● PCBs
Appleton to Little Rapids	○ PCBs	● lead; mercury; PCBs	○ PCBs	○ PCBs	NA	○ mercury; PCBs	● PCBs ○ mercury	● PCBs
Little Rapids to De Pere	● mercury	● lead; mercury; 2,3,7,8-TCDD; PCBs; DDE; DDT	○ mercury; PCBs	○ mercury; PCBs	NA	○ mercury; PCBs	○ mercury; PCBs	● PCBs
Zone 1	○ PCBs	● arsenic; lead; mercury; PCBs; DDD; DDE	○ PCBs; DDE	○ mercury; PCBs; DDE	○ PCBs	○ mercury; PCBs; dieldrin; DDE	○ mercury; PCBs; DDE	● PCBs
Zone 2	● mercury	● mercury; PCBs			○ PCBs; DDE			
Zone 3A		● PCBs	○ PCBs	○ PCBs	NA	○ mercury; PCBs	● PCBs ○ dieldrin	○ PCBs ○ dieldrin
Zone 3B		● arsenic; lead; mercury; PCBs		● PCBs ○ mercury; DDE	NA	● PCBs ○ mercury; dieldrin; DDE	○ mercury; PCBs; DDE	○ dieldrin
Zone 4		● PCBs	NA	○ PCBs; DDE	NA	○ mercury; PCBs	○ mercury; PCBs; DDE	● PCBs

Notes:

NA - No data available.

Risk conclusions based on HQs:

○ - No risk

● - Risk

○ - Potential Risk

Risk conclusions based on weight of evidence:

○ - Site-specific receptor data suggest that there is no risk.

○ - Because of the Federal listing of the bald eagle as threatened, it is concluded that potential risk is actual risk.

and 3B), and carnivorous birds (Green Bay zones 1, 2, 3B, and 4).

(DDT and metabolites) present HQs that were greater than 1.0 for selected receptors.

Sediment Quality Thresholds (SQTs)

SQTs are sediment concentrations that have been linked to a specific magnitude of risk. SQTs were estimated for PCBs with the assumption that a remedy that reduces PCB exposure would also address the other co-occurring COCs. Risk-based concentrations in fish for human and ecological receptors were determined based on:

- Human health cancer risk levels of 10^{-4} , 10^{-5} , and 10^{-6} , and a noncancer hazard index of 1.0 for risk in recreational anglers and high-intake fish consumers
- The NOAECs and LOAECs for species of benthic invertebrates, fish, birds, and riverine mammals found in the river and bay.

SQTs were developed for each pathway and receptor identified as important in the BLRA by the response agencies of the Lower Fox River and Green Bay (e.g., sport fishing consumption, bald eagles). The SQTs themselves are not cleanup criteria, but are used to evaluate levels of PCBs that will be addressed in the Feasibility Study. The final selection of the remedial action levels is a policy decision left to the response agencies. The development and validation of the mathematical model used to define SQTs is described in the BLRA.

To evaluate how PCBs in sediment result in risk to human or ecological receptors, a methodology is needed for translating concentrations of PCBs in sediment to concentrations in fish and higher order organisms. The Fox River Bioaccumulation Model (FRFood Model) was developed for this purpose. FRFood is a series of

mathematical equations that describes a food web and the transfer of bioaccumulating contaminants within that food web. The model includes uptake routes from sediment and water to benthic infauna and ultimately fish, and the model was constructed so that it could be used to either predict fish tissue concentrations from a given sediment concentration, or to predict sediment concentrations from a given fish tissue concentration. The model was validated by running the model “forward;” that is, fish tissue concentrations were predicted from existing sediment concentrations and then compared to measured fish tissue concentrations. When the predicted concentrations were compared to the actual measured concentrations of total PCBs in fish collected in the Lower Fox River and Green Bay, the results were highly comparable.

Estimated SQTs for human health and ecological exposures are shown on Figure 7.

Human Health SQTs

To determine SQTs associated with the protection of human health, fish consumption limits were derived using several different assumptions and risk thresholds. Risk-based fish concentrations (RBFCs) were calculated for recreational anglers and high-intake fish consumers. For recreational anglers, RBFCs were calculated using the average fish intake assumptions from two studies on Michigan anglers (West *et al.*, 1989; West *et al.*, 1993). For high-intake fish consumers, RBFCs were calculated using the average fish intake assumptions for low-income minorities (West *et al.*, 1993) and Hmong (Hutchinson and Kraft, 1994). The RBFCs were generated for each of these exposure scenarios for three

different target risk levels (10^{-6} , 10^{-5} , and 10^{-4}) and for a target noncancer hazard index of 1.0. The RBFCs were used with the results of the FRFood Model to generate a range of SQTs.

Deriving SQTs for each of the consumption scenarios and each of the risks and hazard indices resulted in a total of 48 human health

Ecological SQTs

SQTs protective of ecological receptors were calculated for the Lower Fox River and Green Bay separately. Although the remedial methods may differ between reaches of the river evaluated, the SQTs derived for the De Pere to Green Bay Reach will be applied

Figure 4 Selected PCB HQs that Exceed 1.0





Figure 4 Selected PCB HQs that Exceed 1.0 (Continued)



SQTs, a minimum SQT of $1.1 \mu\text{g/kg}$ (carp at a risk level of 10^{-6} , RME for a high-intake fish consumer) and a maximum SQT of $6,770 \mu\text{g/kg}$ (yellow perch at a risk level of 10^{-4} , CTE for a recreational angler).

to the entire river. These SQTs are based upon levels of total PCBs in fish that either cause risk to the fish themselves, or to birds or mammals that are eating the fish. The SQTs for no observed adverse effects (NOAEC) to walleye is 176, and for carp is

363. The only calculated SQTs that were lower than these for any of the other receptors were the SQT for benthic invertebrates and the SQTs for piscivorous

mammals (mink). The benthic invertebrates threshold effect level (TEL) is a sediment PCB concentration of 31.6 $\mu\text{g/kg}$ and the NOAEC SQT for mink is 24 $\mu\text{g/kg}$.

Figure 5 Selected Mercury HQs that Exceed 1.0 *




Figure 6 Selected DDT or Metabolite HQs that Exceed 1.0





Figure 7. Summary of Sediment Quality Thresholds (SQTs - $\mu\text{g}/\text{kg}$)



Note: Remedial long-term monitoring will include the ecological species listed.

[THIS PAGE LEFT INTENTIONALLY BLANK.]

Table of Contents

1 Introduction	1-1
1.1 Statement of the Problem	1-3
1.2 Objectives of the Baseline Risk Assessment	1-5
1.3 Geographic Boundaries of the Baseline Risk Assessment	1-6
1.4 Organization of the Baseline Risk Assessment	1-7
1.5 Section 1 Figures	1-8
2 Remedial Investigation Summary	2-1
2.1 Environmental Setting and Background	2-1
2.1.1 Site History	2-1
2.1.2 Chemicals of Potential Concern in the Lower Fox River	2-2
2.2 Physical Characteristics	2-4
2.2.1 Total Organic Carbon	2-4
2.2.2 Lower Fox River	2-5
2.2.3 Green Bay	2-6
2.3 Nature and Extent of Contaminants of Potential Concern	2-8
2.3.1 Estimation of PCB Distributions	2-8
2.3.2 Extent of PCB Chemical Impacts	2-12
2.3.3 Extent of Other COC Impacts	2-16
2.4 Contaminant Fate and Transport	2-18
2.4.1 Lower Fox River Sediment Deposition	2-18
2.4.2 Green Bay Sediment Deposition	2-20
2.4.3 PCB Transport	2-21
2.5 Ecological Characteristics (Habitats and Species)	2-22
2.5.1 Open Lands	2-24
2.5.2 Woodlands	2-25
2.5.3 Wetlands	2-26
2.5.4 Riverine Habitat of the Lower Fox River	2-32
2.5.5 Lacustrine Habitat of Green Bay	2-33
2.5.6 Benthic Communities	2-36
2.5.7 Fish	2-38
2.5.8 Birds	2-58
2.5.9 Mink	2-68
2.5.10 Otter	2-71
2.6 Time Trends of Contaminants in Sediment and Fish	2-72
2.6.1 Sediment Methods	2-72
2.6.2 Fish Methods	2-74
2.6.3 Results	2-75
2.6.4 Conclusion	2-80

Table of Contents

2.7	Section 2 Figures and Tables	2-82
3	Summary of the Screening Level Risk Assessment	3-1
3.1	Human Health Screening Level Risk Assessment	3-1
3.1.1	Potential Pathways at Risk	3-1
3.1.2	Chemicals of Potential Concern	3-2
3.2	Ecological Screening Level Risk Assessment	3-2
3.2.1	Potential Pathways at Risk	3-2
3.2.2	Chemicals of Potential Concern	3-3
3.3	Chemicals of Potential Concern for the Baseline Risk Assessment ..	3-3
4	Sediment, Water, and Tissue Chemistry Data	4-1
4.1	Section 4 Figure and Tables	4-2
5	Human Health Risk Assessment	5-1
5.1	Overview	5-1
5.2	Sources, Migration Routes, Human Receptors, and Exposure Pathways	5-3
5.3	Evaluation of Polynuclear Aromatic Hydrocarbons (PAHs) in Whole Body Fish Tissue Samples	5-8
5.3.1	Screening Evaluation	5-9
5.3.2	Calculation of Cancer Risks	5-10
5.3.3	Results of PAH Evaluation	5-11
5.4	Intake Assumptions for Potential Receptors	5-11
5.4.1	Overview of Intake Assumptions	5-12
5.4.2	Generalized Assumptions for Exposure Analysis	5-13
5.4.3	Specific Intake Assumptions for Receptors	5-21
5.5	Exposure Point Concentrations	5-41
5.5.1	Determination of Exposure Point Concentrations	5-42
5.5.2	Source Concentrations	5-44
5.5.3	Transfer Factors and Exposure Point Concentrations	5-49
5.6	Dose-response Assessment	5-49
5.6.1	Overview	5-49
5.6.2	Polychlorinated Biphenyls	5-52
5.6.3	Polychlorinated Dibenzo-p-dioxins and Dibenzofurans	5-68
5.6.4	Dieldrin	5-70
5.6.5	DDT, DDE, and DDD	5-71
5.6.6	Arsenic	5-72

Table of Contents

5.6.7	Lead	5-74
5.6.8	Mercury	5-76
5.6.9	Summary of Toxicity Criteria	5-77
5.7	Baseline Risk Characterization	5-79
5.7.1	Overview	5-79
5.7.2	Recreational Angler	5-81
5.7.3	High-intake Fish Consumer	5-83
5.7.4	Hunter	5-85
5.7.5	Drinking Water User	5-87
5.7.6	Local Resident	5-90
5.7.7	Recreational Water User	5-92
5.7.8	Marine Construction Worker	5-95
5.7.9	Summary of Cancer Risks and Hazard Indices	5-96
5.8	Evaluation of Lead	5-100
5.8.1	Sediment	5-100
5.8.2	Surface Water	5-101
5.8.3	Fish Tissue	5-101
5.8.4	Waterfowl Tissue	5-101
5.9	Focused Evaluation of Exposures to PCBs from Fish Ingestion	5-102
5.9.1	Concentrations of Total PCBs in Fish	5-103
5.9.2	Equations for Calculating Cancer Risks, Hazard Indices, and Target Concentrations in Fish	5-105
5.9.3	Intake Assumptions for Recreational Anglers and High- intake Fish Consumers and Toxicological Parameters	5-107
5.9.4	Cancer Risks and Hazard Indices for Recreational Anglers and High-intake Fish Consumers	5-109
5.9.5	Cancer Risks and Hazard Indices Associated with Intake Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-117
5.9.6	Summary of Probabilistic Risk Assessment	5-120
5.9.7	Evaluation of Exponent Risk Assessment	5-122
5.9.8	Evaluation of PCB Exposures to Young Children	5-124
5.9.9	Risk-based Concentrations in Fish	5-130
5.10	Uncertainty Analysis	5-132
5.11	Summary and Conclusions	5-135
5.11.1	Summary	5-135
5.11.2	Conclusions	5-146
5.12	Section 5 Figures and Tables	5-149

Table of Contents

6 Ecological Risk Assessment	6-1
6.1 Introduction	6-1
6.2 Problem Formulation	6-2
6.2.1 Contaminants of Potential Concern	6-3
6.2.2 Contaminant Fate and Transport	6-3
6.2.3 Assessment and Measurement Endpoints	6-11
6.2.4 Conceptual Site Model	6-18
6.2.5 Food Chain Model	6-29
6.3 Characterization of Ecological Effects	6-37
6.3.1 Data Sources Reviewed	6-37
6.3.2 Polychlorinated Biphenyls (PCBs)	6-40
6.3.3 Dioxins and PCB Congeners	6-62
6.3.4 Dichlorodiphenyl Trichloroethane (DDT)	6-67
6.3.5 Dieldrin	6-73
6.3.6 Arsenic	6-78
6.3.7 Lead	6-80
6.3.8 Mercury	6-80
6.4 Characterization of Exposure	6-89
6.4.1 Exposure Concentrations by Area and Media	6-93
6.4.2 Summary of Exposure Concentrations by Media	6-128
6.4.3 Summary of Exposure Concentrations by Area	6-135
6.5 Risk Characterization	6-144
6.5.1 Estimation of Exposure Point Risks	6-146
6.5.2 Risk Summary by Media	6-157
6.5.3 Risk Summary by Area	6-159
6.5.4 Field Studies in the Lower Fox River and Green Bay	6-180
6.6 Uncertainty	6-195
6.6.1 Uncertainty in the Conceptual Site Model	6-197
6.6.2 Uncertainty in the Data Supporting the Ecological Risk Assessment	6-198
6.6.3 Temporal Variability	6-200
6.6.4 Spatial Variability	6-201
6.6.5 Toxic Exposure Uncertainty	6-203
6.6.6 Alternative Exposure Point Uncertainty	6-205
6.6.7 Population Uncertainty	6-206
6.6.8 Quantitative Uncertainty	6-207
6.7 Risk Management Integration of the Ecological Risk Assessment	6-210

Table of Contents

6.8	Section 6 Figures and Tables	6-231
7	Sediment Quality Thresholds	7-1
7.1	Food Web Models for the Lower Fox River and Green Bay	7-2
7.1.1	FRFood Model	7-3
7.1.2	GBFood Model	7-5
7.2	FRFood Model Food Web Review and Dietary Assignments	7-6
7.3	FRFood Model Calibration	7-7
7.4	Determination of Sediment Quality Thresholds	7-9
7.4.1	Estimating Sediment-to-water Ratios	7-9
7.4.2	Human Health Sediment Quality Thresholds	7-10
7.4.3	Ecological Sediment Quality Thresholds	7-11
7.5	Section 7 Tables	7-12
8	References	8-1

[THIS PAGE LEFT INTENTIONALLY BLANK.]

List of Figures

Figure 1-1	Lower Fox River Study Area	1-9
Figure 1-2	Green Bay Study Area	1-10
Figure 2-1	Lower Fox River Elevation Profile	2-85
Figure 2-2	PCB Distribution (0–10 cm): Little Lake Butte des Morts	2-86
Figure 2-3	PCB Distribution (0–10 cm): Appleton to Little Rapids	2-87
Figure 2-4	PCB Distribution (0–10 cm): Little Rapids to De Pere	2-88
Figure 2-5	PCB Distribution (0–10 cm): De Pere to Green Bay	2-89
Figure 2-6	PCB Distribution (0–10 cm): Green Bay	2-90
Figure 2-7	Estimated Annual Sediment Transport Rates and Stream Flow Velocities	2-91
Figure 2-8	Lower Fox River and Green Bay System Estimated PCB Mass and Major PCB Flux Pathways	2-92
Figure 2-9	Lower Fox River Wetland, Habitat, and Animal Distribution: Little Lake Butte des Morts Reach	2-93
Figure 2-10	Lower Fox River Wetland, Habitat, and Animal Distribution: Appleton to Little Rapids Reach	2-94
Figure 2-11	Lower Fox River Wetland, Habitat, and Animal Distribution: Little Rapids to De Pere Reach	2-95
Figure 2-12	Lower Fox River Wetland, Habitat, and Animal Distribution: De Pere to Green Bay Reach	2-96
Figure 2-13	Wetland Distribution: Green Bay Zones 2 & 3	2-97
Figure 2-14	Wetland Distribution: Green Bay Zone 4	2-98
Figure 2-15	Electrofishing Walleye Catch Data in Green Bay Zone 1	2-99
Figure 2-16	Green Bay Spawning Areas by Fish Types: Salmon/Trout and Benthic Fish	2-100
Figure 2-17	Green Bay Spawning Areas by Fish Types: Pelagic and Game Fish	2-101
Figure 2-18	Green Bay Spawning Areas by Fish Species: Walleye, Yellow Perch, and Sturgeon	2-102
Figure 2-19	Green Bay Spawning Areas by Fish Species: Carp and Alewife . . .	2-103
Figure 2-20	Green Bay Spawning Areas by Fish Species: Emerald Shiners and Gizzard Shad	2-104
Figure 2-21	Distribution of Birds in Green Bay: Select Species and Groups . .	2-105
Figure 2-22	Forster's Tern Population Data in Green Bay	2-106
Figure 2-23	Number of Double-crested Cormorant Nests in Areas 2 and 3 of Green Bay	2-107
Figure 2-24	Distribution of Birds in Green Bay: Eagle and Osprey Locations .	2-108
Figure 2-25	Number of Occupied Bald Eagle Nesting Sites on Green Bay	2-109

List of Figures

Figure 2-26	Mean Annual Productivity of Bald Eagles Nesting on Green Bay, Inland Michigan, and Inland Wisconsin	2-110
Figure 2-27	Lower Fox River Mink Habitat Suitability: Little Lake Butte des Morts Reach	2-111
Figure 2-28	Lower Fox River Mink Habitat Suitability: Appleton to Little Rapids Reach	2-112
Figure 2-29	Lower Fox River Mink Habitat Suitability: Little Rapids to De Pere Reach	2-113
Figure 2-30	Lower Fox River Mink Habitat Suitability: De Pere to Green Bay Reach	2-114
Figure 2-31	Green Bay Mink Habitat Suitability: Zone 2	2-115
Figure 2-32	Green Bay Mink Habitat Suitability: Zone 3	2-116
Figure 2-33	Time Trends of PCBs in Sediments for Depths from 0 to 10 cm and from 10 to 30 cm	2-117
Figure 2-34	Time Trends of PCBs in Sediments for Depths from 30 to 50 cm and from 50 to 100 cm	2-118
Figure 2-35	Time Trends of PCBs in Sediments for Depths over 100 cm	2-119
Figure 4-1	Quality Assurance Status of the Fox River Database Total PCB Data	4-3
Figure 5-1	Potential Source Media, Chemical Migration Routes, Human Receptors, and Exposure Pathways	5-155
Figure 5-2	Cancer Risks for the Little Lake Butte des Morts Reach	5-156
Figure 5-3	Hazard Indices for the Little Lake Butte des Morts Reach	5-157
Figure 5-4	Cancer Risks for the Appleton to Little Rapids Reach	5-158
Figure 5-5	Hazard Indices for the Appleton to Little Rapids Reach	5-159
Figure 5-6	Cancer Risks for the Little Rapids to De Pere Reach	5-160
Figure 5-7	Hazard Indices for the Little Rapids to De Pere Reach	5-161
Figure 5-8	Cancer Risks for the De Pere to Green Bay Reach	5-162
Figure 5-9	Hazard Indices for the De Pere Green Bay Reach	5-163
Figure 5-10	Cancer Risks for Green Bay	5-164
Figure 5-11	Hazard Indices for Green Bay	5-165
Figure 5-12	Range of Cancer Risks for Recreational Anglers and High-intake Fish Consumers in the Lower Fox River	5-166
Figure 5-13	Maximum Cancer Risks for Recreational Anglers and High-intake Fish Consumers by Reach in the Lower Fox River	5-167
Figure 5-14	Range of Cancer Risks for Recreational Anglers and High-intake Fish Consumers in Green Bay	5-168

List of Figures

Figure 5-15	Maximum Cancer Risks for Recreational Anglers and High-intake Fish Consumers by Zone in Green Bay	5-169
Figure 5-16	Range of Hazard Indices for Recreational Anglers and High-intake Fish Consumers in the Lower Fox River	5-170
Figure 5-17	Maximum Hazard Indices for Recreational Anglers and High-intake Fish Consumers by Reach in the Lower Fox River	5-171
Figure 5-18	Range of Hazard Indices for Recreational Anglers and High-intake Fish Consumers in Green Bay	5-172
Figure 5-19	Maximum Hazard Indices for Recreational Anglers and High-intake Fish Consumers by Zone in Green Bay	5-173
Figure 5-20	Comparison of CTE and RME Risk Values with Distribution Data - Little Lake Butte des Morts Reach	5-174
Figure 5-21	Comparison of CTE and RME Hazard Index Values with Distribution Data - Little Lake Butte des Morts Reach	5-175
Figure 5-22	Comparison of CTE and RME Risk Values with Distribution Data - De Pere to Green Bay Reach	5-176
Figure 5-23	Comparison of CTE and RME Hazard Index Values with Distribution Data - De Pere to Green Bay Reach	5-177
Figure 5-24	Risk Variability Evaluation for Recreational Angler - Little Lake Butte des Morts Reach	5-178
Figure 5-25	Hazard Index Variability Evaluation for Recreational Angler - Little Lake Butte des Morts Reach	5-179
Figure 5-26	Risk Variability Evaluation for Recreational Angler - De Pere to Green Bay Reach	5-180
Figure 5-27	Hazard Index Variability Evaluation for Recreational Angler - De Pere to Green Bay Reach	5-181
Figure 5-28	Risk Variability Evaluation for High-intake Fish Consumer - Little Lake Butte des Morts Reach	5-182
Figure 5-29	Hazard Index Variability Evaluation for High-intake Fish Consumer - Little Lake Butte des Morts Reach	5-183
Figure 5-30	Risk Variability Evaluation for High-intake Fish Consumer - De Pere to Green Bay Reach	5-184
Figure 5-31	Hazard Index Variability Evaluation for High-intake Fish Consumer - De Pere to Green Bay Reach	5-185
Figure 5-32	Risk-based Fish Concentrations for Recreational Anglers	5-186
Figure 5-33	Risk-based Fish Concentrations for High-intake Consumers	5-187
Figure 5-34	Risk-based Fish Concentrations Using Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-188

List of Figures

Figure 6-1	Food Web Model: Lower Fox River - Little Lake Butte des Morts to the De Pere Dam	6-239
Figure 6-2	Food Web Model: Green Bay - Zones 1 and 2	6-240
Figure 6-3	Food Web Model: Green Bay - Zones 3 and 4	6-241
Figure 6-4	Data for Egg Mortality and TCDD-Eq	6-242
Figure 6-5	Unfiltered Mercury Concentrations in Surface Water	6-243
Figure 6-6	Total PCB Concentrations (Filtered + Particulate) in Surface Water	6-244
Figure 6-7	Metal Concentrations in Sediments	6-245
Figure 6-8	Total PCB Concentrations in Sediment	6-246
Figure 6-9	Chlorinated Pesticide Concentrations in Sediment	6-247
Figure 6-10	Mercury Concentrations in Fish	6-248
Figure 6-11	Total PCB Concentrations in Yellow Perch, Carp, and Walleye . .	6-249
Figure 6-12	Total PCB Concentrations in Forage Fish	6-250
Figure 6-13	Dieldrin Concentrations in Fish	6-251
Figure 6-14	p,p'-DDE Concentrations in Fish	6-252
Figure 6-15	Measured Total PCB Concentrations in Birds	6-253
Figure 6-16	Measured Dieldrin Concentrations in Birds	6-254
Figure 6-17	Measured p,p'-DDE Concentrations in Birds	6-255
Figure 6-18	PCB Congener Concentrations in Little Lake Butte des Morts Reach	6-256
Figure 6-19	PCB Congener Concentrations in Appleton to Little Rapids Reach	6-257
Figure 6-20	PCB Congener Concentrations in Little Rapids to De Pere Reach .	6-258
Figure 6-21	PCB Congener Concentrations in Green Bay Zones 1 and 2	6-259
Figure 6-22	PCB Congener Concentrations in Green Bay Zone 3A	6-260
Figure 6-23	PCB Congener Concentrations in Green Bay Zone 3B	6-261
Figure 6-24	PCB Congener Concentrations in Green Bay Zone 4	6-262
Figure 6-25	Metal Concentrations in Little Lake Butte des Morts Reach	6-263
Figure 6-26	Total PCB Concentrations in Little Lake Butte des Morts Reach .	6-264
Figure 6-27	Pesticide Concentrations in Little Lake Butte des Morts Reach . .	6-265
Figure 6-28	Metal Concentrations in Appleton to Little Rapids Reach	6-266
Figure 6-29	Total PCB Concentrations in Appleton to Little Rapids Reach . . .	6-267
Figure 6-30	Pesticide Concentrations in Appleton to Little Rapids Reach	6-268
Figure 6-31	Metal Concentrations in Little Rapids to De Pere Reach	6-269
Figure 6-32	Total PCB Concentrations in Little Rapids to De Pere Reach	6-270
Figure 6-33	Pesticide Concentrations in Little Rapids to De Pere Reach	6-271
Figure 6-34	Metal Concentrations in Green Bay Zone 1	6-272

List of Figures

Figure 6-35	Total PCB Concentrations in Green Bay Zone 1	6-273
Figure 6-36	Pesticide Concentrations in Green Bay Zone 1	6-274
Figure 6-37	Metal Concentrations in Green Bay Zone 2	6-275
Figure 6-38	Total PCB Concentrations in Green Bay Zone 2	6-276
Figure 6-39	Dieldrin Concentrations in Green Bay Zone 2	6-277
Figure 6-40	DDT Concentrations in Green Bay Zone 2	6-278
Figure 6-41	Metal Concentrations in Green Bay Zone 3A	6-279
Figure 6-42	Total PCB Concentrations in Green Bay Zone 3A	6-280
Figure 6-43	Pesticide Concentrations in Green Bay Zone 3A	6-281
Figure 6-44	Metal Concentrations in Green Bay Zone 3B	6-282
Figure 6-45	Total PCB Concentrations in Green Bay Zone 3B	6-283
Figure 6-46	Pesticide Concentrations in Green Bay Zone 3B	6-284
Figure 6-47	Metal Concentrations in Green Bay Zone 4	6-285
Figure 6-48	Total PCB Concentrations in Green Bay Zone 4	6-286
Figure 6-49	Pesticide Concentrations in Green Bay Zone 4	6-287
Figure 6-50	Surface Water HQs that Exceeded 1.0	6-288
Figure 6-51	Surface Water Hazard Quotients Exceeding 1.0	6-289
Figure 6-52	Surface Sediment Hazard Quotients that Exceeded 1.0	6-290
Figure 6-53	Surface Sediment Hazard Quotients that Exceeded 1.0	6-291
Figure 6-54	Surface Sediment Metals Hazard Quotients that Exceeded 1.0	6-292
Figure 6-55	Surface Sediment Total PCB Hazard Quotients that Exceeded 1.0	6-293
Figure 6-56	Surface Sediment DDT Hazard Quotients that Exceeded 1.0	6-294
Figure 6-57	Whole Fish Hazard Quotients that Exceeded 1.0	6-295
Figure 6-58	Whole Fish Hazard Quotients that Exceeded 1.0	6-296
Figure 6-59	Whole Fish Mercury Hazard Quotients that Exceeded 1.0	6-297
Figure 6-60	Whole Fish Total PCB Hazard Quotients that Exceeded 1.0	6-298
Figure 6-61	Whole Fish Total PCB Hazard Quotients that Exceeded 1.0	6-299
Figure 6-62	Whole Fish DDE Hazard Quotients that Exceeded 1.0	6-300
Figure 6-63	Bird Hazard Quotients that Exceeded 1.0	6-301
Figure 6-64	Bird Hazard Quotients that Exceeded 1.0	6-302
Figure 6-65a	Bird Total PCB and TEQ Hazard Quotients that Exceeded 1.0 (Reproduction Endpoint)	6-303
Figure 6-65b	Bird Total PCB and TEQ Hazard Quotients that Exceeded 1.0 (Deformity Endpoint)	6-304
Figure 6-66	Bird Metal and Pesticide Hazard Quotients that Exceeded 1.0	6-305
Figure 6-67	Estimated Piscivorous Bird Hazard Quotients that Exceeded 1.0	6-306
Figure 6-68	Estimated Piscivorous Bird Hazard Quotients that Exceeded 1.0	6-307

List of Figures

Figure 6-69	Estimated Piscivorous Bird Mercury Hazard Quotients that Exceeded 1.0	6-308
Figure 6-70	Estimated Piscivorous Bird Total PCB Hazard Quotients that Exceeded 1.0	6-309
Figure 6-71	Comparison of Measured and Estimated Total PCB Hazard Quotients in Piscivorous Birds	6-310
Figure 6-72	Estimated Piscivorous Bird DDE Hazard Quotients that Exceeded 1.0	6-311
Figure 6-73	Estimated Piscivorous Mammal Hazard Quotients that Exceeded 1.0	6-312
Figure 6-74	Estimated Piscivorous Mammal Hazard Quotients that Exceeded 1.0	6-313
Figure 6-75	Estimated Hazard Quotients Exceeding 1.0 in Piscivorous Mammals	6-314
Figure 6-76	Little Lake Butte des Morts Reach Hazard Quotients that Exceed 1.0	6-315
Figure 6-77	Appleton to Little Rapids Reach Hazard Quotients that Exceed 1.0	6-316
Figure 6-78	Little Rapids to De Pere Reach Hazard Quotients that Exceed 1.0	6-317
Figure 6-79	Green Bay Zone 1 Hazard Quotients that Exceed 1.0	6-318
Figure 6-80a	Green Bay Zone 2 Hazard Quotients that Exceed 1.0 (Part 1)	6-319
Figure 6-80b	Green Bay Zone 2 Hazard Quotients that Exceed 1.0 (Part 2)	6-320
Figure 6-80c	Green Bay Zone 2 Hazard Quotients that Exceed 1.0 (Part 3)	6-321
Figure 6-81	Green Bay Zone 3A Hazard Quotients that Exceed 1.0	6-322
Figure 6-82	Green Bay Zone 3B Hazard Quotients that Exceed 1.0	6-323
Figure 6-83	Green Bay Zone 4 Hazard Quotients that Exceed 1.0	6-324
Figure 6-84	Little Lake Butte des Morts Sediment PCB Frequency Distribution	6-325
Figure 6-85	Appleton to Little Rapids Sediment PCB Frequency Distribution	6-326
Figure 6-86	Little Rapids to De Pere Sediment PCB Frequency Distribution	6-327
Figure 6-87	De Pere to Green Bay (Zone 1) Sediment PCB Frequency Distribution	6-328
Figure 6-88	Green Bay Zone 2 Sediment PCB Frequency Distribution	6-329
Figure 6-89	Green Bay Zone 3A Sediment PCB Frequency Distribution	6-330
Figure 6-90	Green Bay Zone 3B Sediment PCB Frequency Distribution	6-331
Figure 6-91	Green Bay Zone 4 Sediment PCB Frequency Distribution	6-332

List of Tables

Table 2-1	Reach and Contaminant Deposit Designations for the Lower Fox River	2-121
Table 2-2	Zone Designations for Green Bay	2-122
Table 2-3	Major Green Bay Wetland Areas/Complexes	2-123
Table 2-4	Green Bay West Shore Wildlife Area Units	2-124
Table 2-5	Summary of Green Bay Tributaries	2-125
Table 2-6	Lower Fox River Habitats	2-126
Table 2-7	Lower Fox River Shoreline and Substrate Types	2-127
Table 2-8	Lower Fox River/Duck Creek Fish Surveys	2-128
Table 2-9	Lower Fox River Fish Species Composition	2-129
Table 2-10	Lower Fox River Fish Populations in the De Pere to Green Bay Reach	2-131
Table 2-11	Green Bay - Common and Important Fish Species	2-133
Table 2-12	Lower Fox River and Green Bay - Common and Important Bird Species	2-134
Table 2-13	Productivity (large young raised per active nest) of Fox River Bald Eagles from 1988 to 1998	2-136
Table 2-14	Endangered and Threatened Mammal Species of the Lower Fox River and Green Bay	2-137
Table 2-15	Results of Sediment Time Trends Analysis for the Lower Fox River	2-138
Table 2-16	Mass-weighted Combined Time Trend for 0 to 10 cm Depth by Reach	2-141
Table 2-17	Results of Fish Time Trends Analysis on the Lower Fox River . . .	2-142
Table 4-1	Description of All Records Contained in the Fox River Database . . .	4-5
Table 4-2	Number of Records Evaluated as Part of the Baseline Risk Assessment - All COPCs	4-6
Table 4-3	Distribution of Existing Sediment, Water, and Tissue Data in the Lower Fox River and Green Bay over Time - Total PCBs Only	4-8
Table 4-4	Distribution of Resident Tissue Samples over Time in the Lower Fox River - Total PCBs Only	4-9
Table 4-5	Distribution of Resident Tissue Samples over Time in Green Bay - Total PCBs Only	4-10
Table 5-1	Potential Human Receptors and Exposure Pathways for the Lower Fox River and Green Bay	5-189
Table 5-2	Fish Consumption Advisories for Lower Fox River and Green Bay	5-190
Table 5-3	Data Summary for 1998 Whole Body Fish Tissue Samples	5-192
Table 5-4	Toxicity Criteria and Calculated RBSCs	5-193

List of Tables

Table 5-5	Screening of Constituents Against RBSCs and Calculated Cancer Risks	5-194
Table 5-6	Permeability Coefficients for Chemicals of Potential Concern . . .	5-195
Table 5-7	Calculated Permeability Coefficients for PCB Aroclors and PCB, Dioxin, and Furan Congeners	5-196
Table 5-8	Absorption Factors for Chemicals for Ingestion of Sediment	5-197
Table 5-9	Absorption Factors for Chemicals for Dermal Contact with Sediment	5-198
Table 5-10	Fish Ingestion Assumptions for Recreational Angler	5-199
Table 5-11	Fish Ingestion Assumptions for High-intake Fish Consumer	5-200
Table 5-12	Consumption of Sport Fish by Hmong Anglers	5-201
Table 5-13	Consumption of Fish from De Pere to Green Bay Reach of Lower Fox River by Hmong/Laotian Anglers	5-202
Table 5-14	Average Size of Meal Consumed by Hmong	5-203
Table 5-15	Summary of Intake Parameter Values for Recreational Anglers—RME Assumptions	5-204
Table 5-16	Summary of Intake Parameter Values for Recreational Anglers—CTE Assumptions	5-205
Table 5-17	Summary of Intake Parameter Values for High-intake Fish Consumers—RME Assumptions	5-206
Table 5-18	Summary of Intake Parameter Values for High-intake Fish Consumers—CTE Assumptions	5-207
Table 5-19	Summary of Intake Parameter Values for Hunters—RME Assumptions	5-208
Table 5-20	Summary of Intake Parameter Values for Hunters—CTE Assumptions	5-209
Table 5-21	Summary of Intake Parameter Values for Drinking Water Users .	5-210
Table 5-22	Summary of Intake Parameter Values for Local Residents	5-211
Table 5-23	Summary of Intake Parameter Values for Swimmers	5-212
Table 5-24	Summary of Intake Parameter Values for Waders	5-213
Table 5-25	Summary of Intake Parameter Values for Marine Construction Workers	5-214
Table 5-26	Upper-bound Measured Concentrations for the Little Lake Butte des Morts Reach	5-215
Table 5-27	Upper-bound Measured Concentrations for the Appleton to Little Rapids Reach	5-216
Table 5-28	Upper-bound Measured Concentrations for the Little Rapids to De Pere Reach	5-217

List of Tables

Table 5-29	Upper-bound Measured Concentrations for the De Pere to Green Bay Reach	5-218
Table 5-30	Upper-bound Measured Concentrations for Green Bay	5-219
Table 5-31	Average Measured Concentrations for the Little Lake Butte des Morts Reach	5-220
Table 5-32	Average Measured Concentrations for the Appleton to Little Rapids Reach	5-221
Table 5-33	Average Measured Concentrations for the Little Rapids to De Pere Reach	5-222
Table 5-34	Average Measured Concentrations for the De Pere to Green Bay Reach	5-223
Table 5-35	Average Measured Concentrations for Green Bay	5-224
Table 5-36	Cancer Evidence for Exposure to Commercial PCB Mixtures	5-225
Table 5-37	PCB Cancer Slope Factors by Persistence and Route of Exposure .	5-226
Table 5-38	Toxicity Equivalency Factors for Dioxin-like PCBs	5-227
Table 5-39	Summary of Dioxin and Furan Toxicity Equivalency Factors	5-228
Table 5-40	Summary of Cancer Slope Factors by Route of Exposure	5-229
Table 5-41	Summary of Reference Doses by Route of Exposure	5-230
Table 5-42	Total Cancer Risks for the Recreational Angler (RME with Upper-bound Concentrations)	5-231
Table 5-43	Total Hazard Indices for the Recreational Angler (RME with Upper-bound Concentrations)	5-232
Table 5-44	Total Cancer Risks for the Recreational Angler (RME with Average Concentrations)	5-233
Table 5-45	Total Hazard Indices for the Recreational Angler (RME with Average Concentrations)	5-234
Table 5-46	Total Cancer Risks for the Recreational Angler (CTE with Average Concentrations)	5-235
Table 5-47	Total Hazard Indices for the Recreational Angler (CTE with Average Concentrations)	5-236
Table 5-48	Total Cancer Risks for the High-intake Fish Consumer (RME with Upper-bound Concentrations)	5-237
Table 5-49	Total Hazard Indices for the High-intake Fish Consumer (RME with Upper-bound Concentrations)	5-238
Table 5-50	Total Cancer Risks for the High-intake Fish Consumer (RME with Average Concentrations)	5-239
Table 5-51	Total Hazard Indices for the High-intake Fish Consumer (RME with Average Concentrations)	5-240

List of Tables

Table 5-52	Total Cancer Risks for the High-intake Fish Consumer (CTE with Average Concentrations)	5-241
Table 5-53	Total Hazard Indices for the High-intake Fish Consumer (CTE with Average Concentrations)	5-242
Table 5-54	Total Cancer Risks for the Hunter (RME with Upper-bound Concentrations)	5-243
Table 5-55	Total Hazard Indices for the Hunter (RME with Upper-bound Concentrations)	5-244
Table 5-56	Total Cancer Risks for the Hunter (RME with Average Concentrations)	5-245
Table 5-57	Total Hazard Indices for the Hunter (RME with Average Concentrations)	5-246
Table 5-58	Total Cancer Risks for the Hunter (CTE with Average Concentrations)	5-247
Table 5-59	Total Hazard Indices for the Hunter (CTE with Average Concentrations)	5-248
Table 5-60	Total Cancer Risks for the Drinking Water User (RME with Upper-bound Concentrations)	5-249
Table 5-61	Total Hazard Indices for the Drinking Water User (RME with Upper-bound Concentrations)	5-250
Table 5-62	Total Hazard Indices for the Drinking Water User (RME with Upper-bound Concentrations and Recent Mercury Data)	5-251
Table 5-63	Total Cancer Risks for the Local Resident (RME with Upper-bound Concentrations)	5-252
Table 5-64	Total Hazard Indices for the Local Resident (RME with Upper-bound Concentrations)	5-253
Table 5-65	Total Hazard Indices for the Local Resident (RME with Upper-bound Concentrations and Recent Mercury Data)	5-254
Table 5-66	Total Cancer Risks for the Recreational Water User: Swimmer (RME with Upper-bound Concentrations)	5-255
Table 5-67	Total Hazard Indices for the Recreational Water User: Swimmer (RME with Upper-bound Concentrations)	5-256
Table 5-68	Total Cancer Risks for the Recreational Water User: Wader (RME with Upper-bound Concentrations)	5-257
Table 5-69	Total Hazard Indices for the Recreational Water User: Wader (RME with Upper-bound Concentrations)	5-258
Table 5-70	Total Cancer Risks for the Marine Construction Worker (RME with Upper-bound Concentrations)	5-259

List of Tables

Table 5-71	Total Hazard Indices for the Marine Construction Worker (RME with Upper-bound Concentrations)	5-260
Table 5-72	Cancer Risks for the Lower Fox River and Green Bay	5-261
Table 5-73	Hazard Indices for the Lower Fox River and Green Bay	5-262
Table 5-74	Summary of Lead Data in Surface Sediment Samples	5-263
Table 5-75	Fish Species with Fillet and Skin Tissue Samples for Total PCBs .	5-264
Table 5-76	Summary of Total PCB Concentrations in Fish Tissue Samples from the Lower Fox River	5-265
Table 5-77	Calculation of PCB Concentration in Carp Fillet Using Fillet-to-Whole Body Ratio	5-266
Table 5-78	Summary of Total PCB Concentrations in Fish Tissue Samples from Green Bay	5-267
Table 5-79	PCB Concentrations in Skin-on Fillet Fish Samples from Lake Winnebago	5-268
Table 5-80	Intake Assumptions and Toxicological Parameters for the Recreational Angler	5-269
Table 5-81	Intake Assumptions and Toxicological Parameters for the High-intake Fish Consumer	5-270
Table 5-82	Cancer Risks by Lower Fox River Reach for the Recreational Angler	5-271
Table 5-83	Cancer Risks by Green Bay Zone for the Recreational Angler . . .	5-272
Table 5-84	Hazard Indices by Lower Fox River Reach for the Recreational Angler	5-273
Table 5-85	Hazard Indices by Green Bay Zone for the Recreational Angler . .	5-274
Table 5-86	Cancer Risks by Lower Fox River Reach for the High-intake Fish Consumer	5-275
Table 5-87	Cancer Risks by Green Bay Zone for the High-intake Fish Consumer	5-276
Table 5-88	Hazard Indices by Lower Fox River Reach for the High-intake Fish Consumer	5-277
Table 5-89	Hazard Indices by Green Bay Zone for the High-intake Fish Consumer	5-278
Table 5-90	Intake Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-279
Table 5-91	Cancer Risks by Lower Fox River Reach Using Intake Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-280

List of Tables

Table 5-92	Cancer Risks by Green Bay Zone Using Intake Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-281
Table 5-93	Hazard Indices by Lower Fox River Reach Using Intake Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-282
Table 5-94	Hazard Indices by Green Bay Zone Using Intake Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-283
Table 5-95	Results of Probabilistic Analysis for Recreational Anglers Using Little Lake Butte des Morts Fish Concentrations	5-284
Table 5-96	Results of Probabilistic Analysis for High-intake Fish Consumers Using Little Lake Butte des Morts Fish Concentrations	5-285
Table 5-97	Results of Probabilistic Analysis for Recreational Anglers Using De Pere to Green Bay Fish Concentrations	5-286
Table 5-98	Results of Probabilistic Analysis for High-intake Fish Consumers Using De Pere to Green Bay Fish Concentrations	5-287
Table 5-99	Summary of Uncertainty Evaluation—Little Lake Butte des Morts Reach	5-288
Table 5-100	Summary of Uncertainty Evaluation—De Pere to Green Bay Reach	5-289
Table 5-101	Child-to-Adult Fish Ingestion Rate Ratios	5-290
Table 5-102	Intake Assumptions and Toxicological Parameters for the Recreational Angler Child	5-291
Table 5-103	Intake Assumptions and Toxicological Parameters for the High-intake Fish Consumer Child	5-292
Table 5-104	Hazard Indices by Lower Fox River Reach for the Recreational Angler Child	5-293
Table 5-105	Hazard Indices by Lower Fox River Reach for the High-intake Fish Consumer Child	5-294
Table 5-106	Risk-based Fish Concentrations for the Recreational Angler	5-295
Table 5-107	Risk-based Fish Concentrations for the High-intake Fish Consumer	5-296
Table 5-108	Risk-based Fish Concentrations Using Intake Assumptions from the Great Lakes Sport Fish Advisory Task Force	5-297
Table 5-109	Cancer Risks for the Lower Fox River and Green Bay	5-298
Table 5-110	Noncancer Hazard Indices for the Lower Fox River and Green Bay	5-299
Table 5-111	Summary of Cancer Risks and Noncancer Hazard Indices for Anglers Exposed to PCBs from Ingestion of Fish	5-300

List of Tables

Table 5-112	Summary of Maximum Cancer Risks and Noncancer Hazard Indices for Anglers Exposed to PCBs from Ingestion of Fish	5-301
Table 5-113	Risk-based Fish Concentrations	5-302
Table 6-1	Fate and Transport Properties of Potentially Bioaccumulating Chemicals of Concern	6-333
Table 6-2	Assessment and Measurement Endpoints for the Ecological Risk Assessment	6-334
Table 6-3	Potential Ecotoxicological Effects from Chemicals Identified in the Lower Fox River/Green Bay	6-335
Table 6-4	Exposure Modeling Input Parameters for Selected Receptor Species	6-336
Table 6-5	Selected Values as Criteria or TRVs	6-337
Table 6-6	Toxic Equivalent Factors (TEFs) Used for TCDD-Eq Calculations	6-340
Table 6-7	Determination of Effects-based TRV for Piscivorous Bird Eggs of the Lower Fox River and Green Bay	6-341
Table 6-8	Surface Water Concentrations in Little Lake Butte des Morts Reach	6-342
Table 6-9	Surface Water Concentrations in Lake Winnebago	6-343
Table 6-10	Surface Sediment Concentrations in Little Lake Butte des Morts Reach	6-344
Table 6-11	Surface Sediment Concentrations in Lake Winnebago	6-345
Table 6-12	PCB Congener Concentrations in Surface Sediment in Little Lake Butte des Morts Reach	6-346
Table 6-13	PCB Congener Concentrations in Surface Sediment in Lake Winnebago	6-347
Table 6-14	Little Lake Butte des Morts Reach Whole Fish Concentrations . .	6-348
Table 6-15	Dioxin and PCB Congener Concentrations in Little Lake Butte des Morts Reach Whole Fish	6-349
Table 6-16	Little Lake Butte des Morts Reach Bird Tissue Concentrations . .	6-351
Table 6-17	Estimated Exposure Concentrations for Piscivorous Birds in Little Lake Butte des Morts Reach	6-352
Table 6-18	PCB Congeners in Tree Swallows from Little Lake Butte des Morts Reach	6-354
Table 6-19	Estimated Exposure Concentrations for Mink in Little Lake Butte des Morts Reach	6-355
Table 6-20	Surface Water Concentrations in Appleton to Little Rapids Reach	6-356
Table 6-21	Surface Sediment Concentrations in Appleton to Little Rapids Reach	6-357

List of Tables

Table 6-22	PCB Congener Concentrations in Surface Sediment in Appleton to Little Rapids Reach	6-358
Table 6-23	Appleton to Little Rapids Reach Whole Fish Concentrations	6-359
Table 6-24	PCB Congener Concentrations in Appleton to Little Rapids Reach Whole Fish	6-360
Table 6-25	Appleton to Little Rapids Reach Bird Tissue Concentrations	6-361
Table 6-26	Estimated Exposure Concentrations for Piscivorous Birds in Appleton to Little Rapids Reach	6-362
Table 6-27	Estimated Exposure Concentrations for Mink in Appleton to Little Rapids Reach	6-364
Table 6-28	Surface Water Concentrations in Little Rapids to De Pere Reach .	6-365
Table 6-29	Surface Sediment Concentrations in Little Rapids to De Pere Reach	6-366
Table 6-30	PCB Congener Concentrations in Surface Sediment in Little Rapids to De Pere Reach	6-367
Table 6-31	Little Rapids to De Pere Reach Whole Fish Concentrations	6-368
Table 6-32	Dioxin and PCB Congener Concentrations in Little Rapids to De Pere Reach Whole Fish	6-369
Table 6-33	Estimated Exposure Concentrations for Piscivorous Birds in Little Rapids to De Pere Reach	6-371
Table 6-34	Estimated Exposure Concentrations for Mink in Little Rapids to De Pere Reach	6-373
Table 6-35	Surface Water Concentrations in De Pere to Green Bay Reach (Green Bay Zone 1)	6-374
Table 6-36	Surface Sediment Concentrations in De Pere to Green Bay Reach (Green Bay Zone 1)	6-375
Table 6-37	PCB Congener Concentrations in Surface Sediment in De Pere to Green Bay Reach (Green Bay Zone 1)	6-376
Table 6-38	De Pere to Green Bay Reach (Green Bay Zone 1) Bird Tissue Concentrations	6-377
Table 6-39	Estimated Exposure Concentrations for Piscivorous Birds in Green Bay Zone 1	6-378
Table 6-40	Estimated Exposure Concentrations for Mink in Green Bay Zone 1	6-380
Table 6-41	Surface Water Concentrations in Green Bay Zone 2 (2A and 2B)	6-381
Table 6-42	Surface Sediment Concentrations in Green Bay Zone 2	6-382
Table 6-43	PCB Congener Concentrations in Surface Sediment in Green Bay Zone 2	6-383

List of Tables

Table 6-44	Green Bay Zones 1 and 2 Whole Fish Concentrations	6-384
Table 6-45	Dioxin and PCB Congener Concentrations in Green Bay Zones 1 and 2 Whole Fish	6-386
Table 6-46	Green Bay Zone 2 Bird Tissue Concentrations	6-388
Table 6-47	Estimated Exposure Concentrations for Piscivorous Birds in Green Bay Zone 2	6-390
Table 6-48	PCB Congeners and Dioxins/Furans in Whole Tree Swallows in Green Bay Zone 2	6-392
Table 6-49	PCB Congeners and Dioxins/Furans in Double-crested Cormorants in Green Bay Zone 2	6-393
Table 6-50	PCB Congeners and Dioxins/Furans in Common Tern Eggs in Green Bay Zone 2	6-395
Table 6-51	PCB Congeners and Dioxins/Furans in Forster's Tern Eggs in Green Bay Zone 2	6-396
Table 6-52	Estimated Exposure Concentrations for Mink in Green Bay Zone 2	6-397
Table 6-53	Surface Water Concentrations in Green Bay Zone 3A	6-398
Table 6-54	Surface Sediment Concentrations in Green Bay Zone 3A	6-399
Table 6-55	PCB Congener Concentrations in Surface Sediment in Green Bay Zone 3A	6-400
Table 6-56	Green Bay Zone 3A Whole Fish Concentrations	6-401
Table 6-57	PCB Congener Concentrations in Green Bay Zone 3A Whole Fish	6-402
Table 6-58	Green Bay Zone 3A Bird Tissue Concentrations	6-403
Table 6-59	Estimated Exposure Concentrations for Piscivorous Birds in Green Bay Zone 3A	6-404
Table 6-60	Estimated Exposure Concentrations for Mink in Green Bay Zone 3A	6-406
Table 6-61	Surface Water Concentrations in Green Bay Zone 3B	6-407
Table 6-62	Surface Sediment Concentrations in Green Bay Zone 3B	6-408
Table 6-63	PCB Congener Concentrations in Surface Sediment in Green Bay Zone 3B	6-409
Table 6-64	Green Bay Zone 3B Whole Fish Concentrations	6-410
Table 6-65	PCB Congener Concentrations in Green Bay Zone 3B Whole Fish	6-411
Table 6-66	Green Bay Zone 3B Bird Tissue Concentrations	6-412
Table 6-67	Estimated Exposure Concentrations for Piscivorous Birds in Green Bay Zone 3B	6-413
Table 6-68	PCB Congeners in Whole Double-crested Cormorants in Green Bay Zone 3B	6-415

List of Tables

Table 6-69	Estimated Exposure Concentrations for Mink in Green Bay Zone 3B	6-416
Table 6-70	Surface Water Concentrations in Green Bay Zone 4	6-417
Table 6-71	Surface Sediment Concentrations in Green Bay Zone 4	6-418
Table 6-72	PCB Congener Concentrations in Surface Sediment in Green Bay Zone 4	6-419
Table 6-73	Green Bay Zone 4 Whole Fish Concentrations	6-420
Table 6-74	PCB Congener Concentrations in Green Bay Zone 4 Whole Fish .	6-421
Table 6-75	Estimated Exposure Concentrations for Piscivorous Birds in Green Bay Zone 4	6-422
Table 6-76	Estimated Exposure Concentrations for Mink in Green Bay Zone 4	6-424
Table 6-77	Hazard Quotients for Whole Fish in Little Lake Butte des Morts Reach	6-425
Table 6-78	PCB Congener and Dioxin/Furan Hazard Quotients for Whole Fish in Little Lake Butte des Morts Reach	6-426
Table 6-79	Hazard Quotients for Bird Tissue in Little Lake Butte des Morts Reach	6-428
Table 6-80	Estimated Hazard Quotients for Piscivorous Birds in Little Lake Butte des Morts Reach	6-429
Table 6-81	PCB Congener Hazard Quotients for Tree Swallows in Little Lake Butte des Morts Reach	6-430
Table 6-82	Estimated Hazard Quotients for Mink in Little Lake Butte des Morts Reach	6-431
Table 6-83	Hazard Quotients for Whole Fish in Appleton to Little Rapids Reach	6-432
Table 6-84	PCB Congener and Dioxin/Furan Hazard Quotients for Whole Fish in Appleton to Little Rapids Reach	6-433
Table 6-85	Hazard Quotients for Bird Tissue in Appleton to Little Rapids Reach	6-434
Table 6-86	Estimated Hazard Quotients for Piscivorous Birds in Appleton to Little Rapids Reach	6-435
Table 6-87	Estimated Hazard Quotients for Mink in Appleton to Little Rapids Reach	6-436
Table 6-88	Hazard Quotients for Whole Fish in Little Rapids to De Pere Reach	6-437
Table 6-89	PCB Congener and Dioxin/Furan Hazard Quotients for Whole Fish in Little Rapids to De Pere Reach	6-438

List of Tables

Table 6-90	Estimated Hazard Quotients for Piscivorous Birds in Little Rapids to De Pere Reach	6-440
Table 6-91	Estimated Hazard Quotients for Mink in Little Rapids to De Pere Reach	6-441
Table 6-92	Hazard Quotients for Bird Tissue in Little Rapids to De Pere Reach	6-442
Table 6-93	Estimated Hazard Quotients for Mink in Green Bay Zone 1	6-443
Table 6-94	Hazard Quotients for Whole Fish in Green Bay Zones 1 and 2 . .	6-444
Table 6-95	PCB Congener and Dioxin/Furan Hazard Quotients for Whole Fish in Green Bay Zones 1 and 2	6-446
Table 6-96	Hazard Quotients for Bird Tissue in Green Bay Zone 2	6-449
Table 6-97	Estimated Hazard Quotients for Piscivorous Birds in Green Bay Zones 1 and 2	6-451
Table 6-98	PCB Congener and Dioxin/Furan Hazard Quotients for Whole Tree Swallows in Green Bay Zone 2	6-452
Table 6-99	PCB Congener and Dioxin/Furan Hazard Quotients for Double-crested Cormorant Eggs and Whole Bodies in Green Bay Zone 2 .	6-453
Table 6-100	PCB Congener and Dioxin/Furan Hazard Quotients for Common Tern Eggs in Green Bay Zone 2	6-454
Table 6-101	PCB Congener and Dioxin/Furan Hazard Quotients for Forster's Tern Eggs in Green Bay Zone 2	6-455
Table 6-102	Estimated Hazard Quotients for Mink in Green Bay Zone 2	6-456
Table 6-103	Hazard Quotients for Whole Fish in Green Bay Zone 3A	6-457
Table 6-104	PCB Congener Hazard Quotients for Whole Fish in Green Bay Zone 3A	6-458
Table 6-105	Hazard Quotients for Bird Tissue in Green Bay Zone 3A	6-459
Table 6-106	Estimated Hazard Quotients for Piscivorous Birds in Green Bay Zone 3A	6-460
Table 6-107	Estimated Hazard Quotients for Mink in Green Bay Zone 3A . . .	6-461
Table 6-108	Hazard Quotients for Whole Fish in Green Bay Zone 3B	6-462
Table 6-109	PCB Congener Hazard Quotients for Whole Fish in Green Bay Zone 3B	6-463
Table 6-110	Hazard Quotients for Bird Tissue in Green Bay Zone 3B	6-464
Table 6-111	Estimated Hazard Quotients for Piscivorous Birds in Green Bay Zone 3B	6-465
Table 6-112	PCB Congener Hazard Quotients for Double-crested Cormorants in Green Bay Zone 3B	6-466
Table 6-113	Estimated Hazard Quotients for Mink in Green Bay Zone 3B . . .	6-467

List of Tables

Table 6-114	Hazard Quotients for Whole Fish in Green Bay Zone 4	6-468
Table 6-115	PCB Congener Hazard Quotients for Whole Fish in Green Bay Zone 4	6-469
Table 6-116	Estimated Hazard Quotients for Piscivorous Birds in Green Bay Zone 4	6-470
Table 6-117	Estimated Hazard Quotients for Mink in Green Bay Zone 4	6-471
Table 6-118	Bald Eagle Productivity in the Lower Fox River and Green Bay (from Dykstra and Meyer, 1996)	6-472
Table 6-119	Summary of Bald Eagle Productivity in the Lower Fox River and Green Bay (from Dykstra and Meyer, 1996)	6-473
Table 6-120	Organochlorine Concentrations in Addled Eggs (from Dykstra and Meyer, 1996)	6-474
Table 6-121	Summary of BLRA Risk Assessment Results for Little Lake Butte des Morts Reach	6-475
Table 6-122	Summary of BLRA Risk Assessment Results for Appleton to Little Rapids Reach	6-477
Table 6-123	Summary of BLRA Risk Assessment Results for Little Rapids to De Pere Reach	6-479
Table 6-124	Summary of BLRA Risk Assessment Results for Green Bay Zone 1	6-481
Table 6-125	Summary of BLRA Risk Assessment Results for Green Bay Zone 2	6-482
Table 6-126	Summary of BLRA Risk Assessment Results for Green Bay Zone 3A	6-484
Table 6-127	Summary of BLRA Risk Assessment Results for Green Bay Zone 3B	6-486
Table 6-128	Summary of BLRA Risk Assessment Results for Green Bay Zone 4	6-487
Table 6-129	Summary of Published Sediment Quality Criteria	6-488
Table 6-130	Hazard Quotients for 90 th Percentile COPC Concentrations in Walleye	6-491
Table 6-131	Hazard Quotients for 90 th Percentile PCB Congener Concentrations in Whole Walleye	6-492
Table 6-132	Hazard Quotients for 90 th Percentile COPC Concentrations in Double-crested Cormorants	6-493
Table 6-133	Hazard Quotients for 90 th Percentile PCB Congener Concentrations in Double-crested Cormorants	6-494
Table 6-134	Ecological Risk Summary Table	6-495

List of Tables

Table 7-1	References Reviewed for Potential Input Parameter to the Lower Fox River Bioaccumulation Model	7-13
Table 7-2	Inputs to the FRFood Model for Model Calibration in Little Lake Butte des Morts Reach	7-14
Table 7-3	Inputs to the FRFood Model for Model Calibration in Little Rapids to De Pere Reach	7-15
Table 7-4	Inputs to the FRFood Model for Model Calibration in Green Bay Zone 1	7-16
Table 7-5	Inputs to the FRFood Model for Model Calibration in Green Bay Zone 2	7-17
Table 7-6	Lower Fox River Bioaccumulation Model Calibration	7-18
Table 7-7	Reach-specific and River-wide Total PCB Water-to-Sediment Ratios	7-20
Table 7-8	Ratio of PCB Concentrations in Fillet to Whole Body for Different Species	7-21
Table 7-9	Sediment Quality Thresholds Estimated for Human Health Effects at a 10^{-5} Cancer Risk and Noncancer Hazard Index of 1.0 .	7-22
Table 7-10	Derivation of Bird Biomagnification Factors (BMFs) for Total PCBs	7-23
Table 7-11	Sediment Quality Thresholds Estimated for Ecological Effects	7-24

[THIS PAGE LEFT INTENTIONALLY BLANK.]

List of Appendices

Appendix A Letter from David Webb, Wisconsin Department of Natural Resources,
August 3, 1998

Appendix B Human Health Fate and Transport Models, Transport Factors, and
Reduction Factors

B1 Additional Evaluation of Exposure to PCBs in Fish from the Lower Fox
River and Green Bay

B2 General Statistics

B3 Fate and Transport Models and Transfer Factors

B4 Exposure Point Concentrations, Unit Cancer Risks, Unit Hazard
Indices, Cancer Risks, and Hazard Indices for Different Receptors

Appendix C Focused Ecological Risk Assessment Upper Green Bay Portion of the
Fox River Site, Green Bay, Wisconsin

[THIS PAGE LEFT INTENTIONALLY BLANK.]

List of Acronyms

2,3,7,8-TCDD	2,3,7,8-tetrachloro- <i>p</i> -dibenzodioxin
2,3,7,8-TCDF	2,3,7,8-tetrachloro- <i>p</i> -dibenzofuran
95% UCL	95 percent upper confidence limit
°C	degrees centigrade
°F	degrees Fahrenheit
μg	microgram
μg/dl	micrograms per deciliter
μg/dl-blood	micrograms per deciliter of blood
μg/kg	micrograms per kilogram
μg/kg-BW/day	micrograms per kilogram of body weight per day
μg/kg-day	micrograms per kilogram per day
μg/kg-fillet	micrograms per kilogram of fish fillet
μg/kg-whole body	micrograms per kilogram of whole-body fish
μg/kg-sediment	micrograms per kilogram of sediment
μg/L	micrograms per liter
μg/m ³	micrograms per cubic meter
μg-PCB/kg-BW/day	micrograms of polychlorinated biphenyl per kilogram of body weight per day
μg-TCDD/kg-lipid	micrograms of 2,3,7,8-tetrachloro- <i>p</i> -dibenzodioxin per kilogram of lipid
μm	micrometer
ABS	ingestion absorption factor (fraction absorbed) or inhalation absorption factor (fraction absorbed)
AChE	acetylcholinesterase
ADD	average daily dose
AE	assimilation efficiency (in %)
AEHS	Association for the Environmental Health of Soils
AF	sediment adherence factor (in mg/cm ²)
a_{f-wb}	ratio of concentrations in fish fillet to concentrations in whole body of fish (in kg-fish/kg-fillets)
AHH	aryl hydrocarbon hydroxylase
Ah-R	aryl hydrocarbon receptor
AQUIRE	Aquatic Information Retrieval Database
ARCS	Assessment and Remediation of Contaminated Sediments
As ³⁺	arsenite (trivalent arsenic compound)
As ⁵⁺	arsenate
AT	averaging time (in days)
ATc	averaging time (carcinogenic)

List of Acronyms

AT_{nc}	averaging time (non-carcinogenic) or averaging time for chronic, noncancer effects (in days)
AT_{nc_C}	non-carcinogenic averaging time for a child
ATSDR	Agency for Toxic Substances and Disease Registry (part of the United States Public Health Service)
BAF	bioaccumulation factor
BLRA	baseline risk assessment
BLRPC	Bay Lake Regional Planning Commission
BMF	biomagnification factor
BSAF	biota-to-sediment accumulation factor
BTAG	Biological Technical Assistance Group
BW or bw	body weight (in kg)
BW_C	body weight for a child
C	chemical concentration (in mg/kg-soil or mg/L- water)
CA	concentration of chemical in air (in mg/m ³)
C_{ab}	chemical concentration in indoor air during a bath
C_{as}	chemical concentration in indoor air during a shower
CDF	confined disposal facility
CERCLA	Comprehensive Environmental Response, Compensation and Liability Act of 1980 (the Superfund statute)
CF	conversion factor (in kg/g or kg/mg) or volumetric conversion factor (in L/1,000 cc)
C_{fish}	chemical concentration in fish (in mg/kg-fish)
$C_{fish_{EPC}}$	exposure point concentration in fish
C_{fish-f}	concentration of PCBs in fish fillet (in µg/kg-fillet)
$C_{fish_{meas}}$	measured fish chemical concentration
$C_{fish_{meas_i}}$	measured concentration of chemical i in fish (in mg/kg)
$C_{fish-wb}$	concentration of PCBs in whole body of fish (in µg/kg-whole body)
cfs	cubic feet per second
cm	centimeter
cm ²	square centimeter
cm ² /event	square centimeters per event
cm/hr	centimeters per hour
C_{oa}	chemical concentration in outdoor air

List of Acronyms

COC	chemical of concern
COPC	chemical of potential concern
C_{pw}	chemical concentration in sediment pore water
CR	contact rate or the amount of impacted medium contacted per event
CS	chemical concentration in sediment (in mg/kg-sediment)
C_{sed}	measured sediment chemical concentration
CSF	cancer slope factor
CSF_d	cancer slope factor for evaluating absorbed dermal doses (in [mg/kg-day] ⁻¹)
CSF_i	inhalation cancer slope factor
CSF_o	cancer slope factor for evaluating administered ingestion doses (in [mg/kg-day] ⁻¹)
CSF_o	oral cancer slope factor (in [mg/kg-day] ⁻¹)
C_{sw}	chemical concentration in surface water
C_{sw-di}	measured dissolved concentration for chemical <i>i</i> in water (in mg/L)
C_{sw-ti}	measured total concentration of chemical <i>i</i> in water (in mg/L)
CTE	central tendency exposure
CW	chemical concentration in water (in mg/L)
C_{wb}	chemical concentration in bath water
CWF	chemical concentration in waterfowl (in mg/kg-waterfowl)
CWF_{EPC}	exposure point concentration in waterfowl
CWF_{meas}	measured chemical concentration in waterfowl
CWF_{measi}	measured concentration of chemical <i>i</i> in waterfowl (in mg/kg)
C_{ws}	chemical concentration in shower water
C_x	concentration of the COPC in medium <i>x</i> (in mg/kg ww)
cy	cubic yard
days/yr	days per year
DDD	4,4'-dichlorodiphenyl dichloroethane (includes isomers o,p'-DDD and p,p'-DDD)
DDE	4,4'-dichlorodiphenyl dichloroethylene (includes isomers o,p'-DDE and p,p'-DDE)
DDOH-PA	metabolite of DDT conjugated to a fatty acid

List of Acronyms

DDT	4,4'-dichlorodiphenyl trichloroethane (includes isomers o,p'-DDT and p,p'-DDT)
DNA	deoxyribonucleic acid
D.O.	dissolved oxygen
dwt	dry weight
EC ₂₀	20 percent effect concentration
EC ₃₀	30 percent effect concentration
EC ₅₀	50 percent effect concentration
ED	exposure duration (in years)
ED _C	exposure duration for a child
ED _T	estimated daily dose (in mg/kg-BW/day ww)
EEC	Exposure Effect Concentration or Extreme Effect Concentration
FRFood	Fox River Food Model
FRG	Fox River Group, which is composed of the following seven companies (listed alphabetically): Appleton Papers, Inc.; Fort James Corporation; NCR Corporation; P. H. Glatfelter Company; Riverside Paper Corporation; U.S. Paper Mills Corporation; and Wisconsin Tissue Mills, Inc.
FS	feasibility study
g	gram
GAS	Graef, Anhalt, Schloemer and Associates, Inc.
GBFood	Green Bay Food Model
GBTOXe	Green Bay Toxics Model
g/day	grams per day
GE	gross energy (in kcal/g)
g-fish/day	grams of fish per day
GLEMEDS	Great Lakes Embryo Mortality, Edema, and Deformities Syndrome
GLWQI	Great Lakes Water Quality Initiative
g/meal	grams per meal
g/mole	grams per mole
g-waterfowl/day	grams of waterfowl per day
g/yr	grams per year
H ⁺	protons
HEAST	Health Effects Assessment Summary Table
Hg ⁰	elemental mercury
Hg ²⁺	mercuric ion

List of Acronyms

Hg_2^{2+}	mercurous ion
HgOH	inorganic mercury
HI	hazard index
Hi_i	hazard index for chemical i
HQ	hazard quotient
hrs/day	hours per day
I	chemical intake (in mg/kg-BW/day)
I_0	interpolated zeroed grid
I_d	interpolated deleted grid
I_{der-s}	absorbed dose from dermal contact with sediment (in mg/kg-BW/day)
I_{der-w}	absorbed intake from dermal contact with water (in mg/kg-BW/day) = $TBS \cdot FBE$
IEUBK	Integrated Exposure Biokinetic/Uptake Model
I_{ing-f}	intake from ingestion of fish (in mg/kg-BW/day)
I_{ing-s}	intake from incidental ingestion of sediment (in mg/kg-BW/day)
I_{ing-w}	intake from ingestion of water (in mg/kg-BW/day)
I_{ing-wf}	intake from ingestion of waterfowl (in mg/kg-BW/day)
I_{inhal}	intake from inhalation (in mg/kg-BW/day)
Inc	intake from ingestion of fish averaged over the exposure period (in mg/kg-day)
$IntFacC$	intake factor for cancer risk (in $[\text{mg/kg}]^{-1}$)
$IntFacNC$	intake factor for chronic, noncancer effects (in $[\text{mg/kg}]^{-1}$)
IPS	Integrated Paper Services
IR	ingestion rate (in g/day or L/day) or inhalation rate (in m^3/hour) or incidental sediment ingestion rate (in mg-sediment/day)
IR_A	ingestion rate for an adult
IR_C	ingestion rate for a child
IRIS	Integrated Risk Information System
IUPAC	International Union of Pure and Applied Chemistry
I_x	rate of ingestion of medium x (in mg/day or kg/day ww)
kcal/day	kilocalories per day
kcal/g	kilocalories per gram

List of Acronyms

kg	kilogram (1 kg is approximately equivalent to 2.2 pounds)
kg-fish/kg-fillets	kilograms of fish-to-kilograms of fillets
kg/g	kilograms per gram
kg/L	kilograms per liter
kg/mg	kilograms per milligram
km	kilometer (1 km is approximately equivalent to 0.6 mile)
km ²	square kilometer
km ³	cubic kilometer
K_{ow}	octanol-water partitioning coefficient
K_p	permeability coefficient
LADD	lifetime average daily dose
LC ₁₀	10 percent lowest effect concentration
LC ₁₂	12 percent lowest effect concentration
LC ₅₀	50 percent lowest effect concentration
LC ₉₀	90 percent lowest effect concentration
L/1,000 cc	liters per 1,000 cubic centimeters
LD ₁₀	lethal dose to 10 percent of test population
LD ₂₀	lethal dose to 20 percent of test population
LD ₃₀	lethal dose to 30 percent of test population
LD ₅₀	lethal dose to 50 percent of test population
L/day	liters per day
LLBdM	Little Lake Butte des Morts
L/m ³	liters per cubic meter
L/mg	liters per milligram
LOAEC	Lowest Observed Adverse Effect Concentration
LOAEL	Lowest Observed Adverse Effect Level
LOEL	Lowest Observed Effect Level
m ²	square meter
m ³	cubic meter
MDNR	Michigan Department of Natural Resources
ME	metabolizable energy (in kcal/g prey)
meals/yr	meals per year
MEC	Moderate Effect Concentration
MeHg	methylmercury (organic mercury)
MFO	mixed function oxidase
mg	milligram

List of Acronyms

mg-Aroclor 1254/kg-BW/day	milligrams of Aroclor 1254 per kilogram of body weight per day
mg/cm ²	milligrams per square centimeter
mg/day	milligrams per day
mg-Hg/kg-BW/day	milligrams of mercury per kilogram of body weight per day
mg/kg	milligrams per kilogram
mg/kg-BW	milligrams per kilogram of body weight
mg/kg-BW/day	milligrams per kilogram of body weight per day
mg/kg-day	milligrams per kilogram per day
mg/kg-DDE	milligrams per kilogram of 4,4'-dichlorodiphenyl dichloroethylene
mg/kg-DDT	milligrams per kilogram of 4,4'-dichlorodiphenyl trichloroethane
mg/kg-egg	milligrams per kilogram of egg
mg/kg-fish	milligrams per kilogram of fish
mg/kg-sediment	milligrams per kilogram of sediment
mg/kg-soil	milligrams per kilogram of soil
mg/kg-waterfowl	milligrams per kilogram of waterfowl
mg/L	milligrams per liter
mg/L-water	milligrams per liter of water
mg/m ³	milligrams per cubic meter
mg/mg	milligrams per milligram
mg-sediment/day	milligrams of sediment per day
mg/yr	milligrams per year
m ³ /hr	cubic meters per hour
mi ²	square mile
mi ³	cubic mile
ml/day	milliliters per day
mm	millimeter
m ³ /mg	cubic meters per milligram
MNFI	Michigan Natural Features Inventory
MRL	Minimal Risk Level
m/s	meters per second
m ³ /s	cubic meters per second
MSA	Metropolitan Statistical Area
MT	metric ton
MW	molecular weight (in g/mole)
N	non-interpolated grid

List of Acronyms

NASS	National Agricultural Statistics Service
NAWQC	National Ambient Water Quality Criteria
NCP	National Contingency Plan
NCR	National Cash Register
“ND”	no data
NEC	No Effect Concentration
ng/kg	nanograms per kilogram
ng/kg-egg	nanograms per kilogram of egg
ng/kg-TCDD/egg	nanograms per kilogram of 2,3,7,8-tetrachloro- <i>p</i> -dibenzodioxin per egg
ng/kg-TEQ/egg	nanograms per kilogram of toxic equivalency per egg
ng/kg-ww/eagle	nanograms per kilogram of wet weight per eagle
ng/kg-ww/egg	nanograms per kilogram of wet weight per egg
ng/L	nanograms per liter
ng-TEQ/kg-ww/egg	nanograms of toxic equivalency per kilogram of wet weight per egg
NOAA	National Oceanic and Atmospheric Administration
NOAEC	No Observed Adverse Effect Concentration
NOAEL	No Observed Adverse Effect Level
NOEL	No Observed Effect Level
NRDA	Natural Resource Damage Assessment
N.W.R.	National Wildlife Refuge
OMOE	Ontario Ministry of the Environment
PAH	polynuclear aromatic hydrocarbon
PC	permeability constant (in cm/hr)
PCB	polychlorinated biphenyl
PCDD	polychlorinated dibenzodioxin
PCDF	polychlorinated dibenzofuran
PCH	planar chlorinated hydrocarbon
PCP	pentachlorophenol
pg	picogram
pg/g	picograms per gram
pg/kg-day	picograms per kilogram per day
PHH	planar halogenated hydrocarbons
ppb	parts per billion
ppm	parts per million
ppt	parts per trillion
PRP	potentially responsible party
QA	quality assurance

List of Acronyms

QA/QC	quality assurance/quality control
<i>R</i>	cancer risk
RA	risk assessment
<i>Ratio</i> _{CAFI}	child-to-adult fish ingestion ratio
RBFC	risk-based fish concentration
RBSC	risk-based screening concentration
<i>RBSC</i> _{SA-fish}	high-intake fish consumer risk-based screening concentration for carcinogenic or non-carcinogenic chemicals
RETEC	Remediation Technologies, Inc.
<i>RF</i>	reduction factor
RfC	EPA Reference Concentration
<i>RfD</i>	chronic oral reference dose (chemical-specific) or EPA Reference Dose
<i>RfD_d</i>	reference dose for evaluating absorbed dermal doses (in mg/kg-day)
<i>RfD_o</i>	reference dose for evaluating administered ingestion doses (in mg/kg-day)
<i>RfDo</i>	oral reference dose for chronic, noncancer effects (in mg/kg-day)
<i>RF_{fish}</i>	reduction factor for fish
<i>RF_{fishi}</i>	reduction factor for chemical <i>i</i> for fish (in mg/mg)
<i>RF_{WF}</i>	reduction factor for waterfowl
<i>RF_{WFi}</i>	reduction factor for chemical <i>i</i> for waterfowl (in mg/mg)
<i>R_i</i>	cancer risk for chemical <i>i</i>
RI	remedial investigation
RI/FS	remedial investigation and feasibility study
RME	reasonable maximum exposure
ROD	Record of Decision
SA	exposed skin surface area (in cm ² or cm ² /event) = <i>TBS</i> · <i>FBE</i>
SAIC	Science Applications International Corporation
SAV	submerged aquatic vegetation and/or floating vegetation
SCS	Soil Conservation Service
SEC	Sediment Effect Concentration
<i>SF</i>	oral cancer slope factor (chemical-specific)
SLRA	screening level risk assessment

List of Acronyms

SMDP	Scientific Management Decision Point
SMU	sediment management unit
SQC	Sediment Quality Criteria
SQT	sediment quality threshold
SVOC	semivolatile organic compound
SWAC	sediment-weighted average concentration
TBS	total body surface area (in cm^2)
TCDD	2,3,7,8-tetrachloro- <i>p</i> -dibenzodioxin
TCDD-Eq	2,3,7,8-tetrachloro- <i>p</i> -dibenzodioxin equivalent
TCDF	2,3,7,8-tetrachloro- <i>p</i> -dibenzofuran
TEC	Threshold Effect Concentration
TEF	toxic equivalency factor
TEL	Environmental Canada Threshold Effect Level
TEQ	toxic equivalency
TF_{bwa}	bath water-to-air transfer factor
Tf_{bwai}	transfer factor for chemical i for volatilization from bath water to air (in L/m^3)
TF_{sdpw}	sediment-to-pore water transfer factor
Tf_{sdpi}	transfer factor for chemical i for sediment to pore water (in kg/L)
TF_{sh}	shower water-to-air transfer factor
Tf_{shi}	transfer factor for chemical i for volatilization from shower water to air (in L/m^3)
TF_{swoa}	surface water-to-air transfer factor
Tf_{swoai}	transfer factor for volatilization from surface water to outdoor air (in L/m^3)
THI	target hazard index
THQ	target hazard quotient
TIE	Toxicity Evaluation Identification
TOC	total organic carbon
TR	target risk
TRV	Toxicity Reference Value
TSS	total suspended solids
$UHIa1\text{-inh-}c_i$	unit hazard index for chemical i for inhalation of outdoor air by a young child (in m^3/mg)
$UHIa2\text{-inh-}c_i$	unit hazard index for chemical i for inhalation of outdoor air (in m^3/mg)
$UHIfd1\text{-ing-}c_i$	unit hazard index for chemical i for ingestion of waterfowl (in kg/mg)

List of Acronyms

$UHI_{fsh1-ing-c_i}$	unit hazard index for chemical i for ingestion of fish (in kg/mg)
$UHI_{sd1-d-c_i}$	unit hazard index for chemical i for dermal contact with sediment (in kg/mg)
$UHI_{sd1-ing-c_i}$	unit hazard index factor for chemical i for ingestion of sediment (in kg/mg)
$UHI_{w1av-inh-c_i}$	unit hazard index for chemical i for inhalation of indoor air by a young child (in m ³ /mg)
UHI_{w1-d-c_i}	unit hazard index for chemical i for dermal contact with surface water by a young child (in L/mg)
$UHI_{w1-ing-c_i}$	unit hazard index for chemical i for incidental ingestion of surface water by a young child (in L/mg)
$UHI_{w2av-inh-c_i}$	unit hazard index for chemical i for inhalation of indoor air by an adult (in m ³ /mg)
UHI_{w2-d-c_i}	unit hazard index for chemical i for dermal contact with surface water (in L/mg)
$UHI_{w2-ing-c_i}$	unit hazard index for chemical i for incidental ingestion of surface water (in L/mg)
UHI_{w3-d-c_i}	unit hazard index for chemical i for dermal contact with sediment pore water (in L/mg)
UP	Michigan's Upper Peninsula
URF	unit risk factor
$URF_{a1-inh-c_i}$	unit risk factor for chemical i for inhalation of outdoor air by a young child (in m ³ /mg)
$URF_{a2-inh-c_i}$	unit risk factor for chemical i for inhalation of outdoor air (in m ³ /mg)
$URF_{fd1-ing-c_i}$	unit risk factor for chemical i for ingestion of waterfowl (in kg/mg)
$URF_{fsh1-ing-c_i}$	unit risk factor for chemical i for ingestion of fish (in kg/mg)
URF_i	inhalation unit risk factor
$URF_{sd1-d-c_i}$	unit risk factor for chemical i for dermal contact with sediment (in kg/mg)
$URF_{sd1-ing-c_i}$	unit risk factor for chemical i for ingestion of sediment (in kg/mg)
$URF_{w1av-inh-c_i}$	unit risk factor for chemical i for inhalation of indoor air by a young child (in m ³ /mg)
URF_{w1-d-c_i}	unit risk factor for chemical i for dermal contact with surface water by a young child (in L/mg)

List of Acronyms

$URFw1-ing-c_i$	unit risk factor for chemical i for incidental ingestion of surface water by a young child (in L/mg)
$URFw2av-inh-c_i$	unit risk factor for chemical i for inhalation of indoor air by an adult (in m ³ /mg)
$URFw2-d-c_i$	unit risk factor for chemical i for dermal contact with surface water (in L/mg)
$URFw2-ing-c_i$	unit risk factor for chemical i for incidental ingestion of surface water (in L/mg)
$URFw3-d-c_i$	unit risk factor for chemical i for dermal contact with sediment pore water (in L/mg)
USACE	United States Army Corps of Engineers
USDA	United States Department of Agriculture
USFWS	United States Fish and Wildlife Service
USGS	United States Geological Survey
UWSGI	University of Wisconsin Sea Grant Institute
W.A.	Wildlife Area
WDH	Wisconsin Department of Health and Social Services
WDNR	Wisconsin Department of Natural Resources
WHO	World Health Organization
wLFRM	Whole Lower Fox River Model
WSEV	Window Subsampling Empirical Variance
ww	wet weight
YOY	young-of-the-year